

First Recovery of *Schistosoma mansoni* Eggs From a Latrine in Europe (15–16th Centuries)

F. Bouchet, S. Harter, J. C. Paicheler, A. Araújo*, and L. F. Ferreira*, Laboratoire de Parasitologie et de Paléoparasitologie, EA 3308, associé CNRS ESA 8045, U.F.R de Pharmacie, Université de Reims, 51, rue Cognacq-Jay, 51096 REIMS Cedex, France; *Fundação Oswaldo Cruz, Escola Nacional de Saúde Pública, Departamento de Endemias Samuel Pessoa, rua Leopoldo Bulhões, 1480, 21041-210, Rio de Janeiro, Brazil. e-mail: francoise.bouchet@univ-reims.fr

ABSTRACT: *Schistosoma mansoni* eggs were found in a latrine dated AD 1450–1550 in Montbéliard, France. This is the first record of intestinal schistosomiasis in archaeological material. Because the parasite has an African origin, its presence in Europe at that time may be due to African slaves. However, intestinal schistosomiasis has never been established in Europe.

Human and other animal intestinal parasite eggs have been recorded from a latrine dating AD 1450–1550 in Montbéliard, France (Bouchet and Paicheler, 1995). Reviewing the same material, helminth eggs with a typical *Schistosoma mansoni* lateral spine were also found.

The material studied was composed of coprolites and sediment collected from a latrine found in a house built during the second half of the 15th century. The latrine (2.35 × 2.30 × 1.65 m) is located in an inner courtyard, which was completely closed from the first half of the 16th century up to the recent archaeological excavations. Three stratigraphic units were identified, but only samples of the middle one

(US544) were sent for paleoparasitological analysis. The sediment is macroscopically rich in organic remains, fecal material, and artifacts. Dishware and other cultural remains point to very rich dwellings (Cantrelle et al., 2000).

The recovered parasite eggs were found to be very well preserved by the anaerobic and humidity conditions. Thus, the material was not rehydrated by current methods used in palaeoparasitology (Reinhard et al., 1988). Samples were directly submitted to ultrasound for 20 sec then run through graduated sieves (315, 160, 50, and 25 μm). The final sediment recovered from the last 2 sieves was observed under a microscope. Eggs and other organic remains found were measured and photographed.

The eggs presented here were identified by their morphology as *S. mansoni*, size (142 × 58 μm), and lateral spine (Fig. 1). All 4 species classified in the *S. mansoni* group have eggs with a lateral spine. Apart from *S. mansoni*, these species are found only in Africa. *Schistosoma rodhaini* is a parasite found in African rodents and carnivores; *Schistosoma edwardiense*, which have eggs with a rudimentary lateral spine, is a parasite of *Hippopotamus amphibius*; *Schistosoma hippopotami*, in the same host, might be a synonym of *S. mansoni* (Rollinson and Simpson, 1987).

The palaeoparasitology of human schistosomiasis has been discussed by several authors. Wei (1973) described a *Schistosoma japonicum* infection in a mummified body of the Han Dynasty. *Schistosoma japonicum* eggs were found in China in a 2,100-yr-old male corpse (Liangbiao and Hung, 1981). *Schistosoma haematobium* was the first parasite found in an Egyptian mummy dated 5,200-yr-old (Ruffer, 1910). Later, as Egyptian mummy necropsies were performed more frequently, this parasite was found at different periods (Jonckheere, 1942; Reyman, 1973; Cockburn et al., 1975; Lewin, 1977; Capron, 1993; Contis and David, 1996). The advent of new techniques allowed the diagnosis of other cases (Deelder et al., 1990; David, 1997), and a paleoepidemiology of schistosome infection was tentatively drawn (Miller et al., 1992; Araújo and Ferreira, 1997).

There are no records of human schistosomiasis infection outside Asian or African prehistoric populations (Araújo and Ferreira, 1997). *Schistosoma haematobium* infection was reported in Europe in the 19th century but vanished in a short period of time (Brumpt, 1936). *Schistosoma haematobium* eggs dated to the 15th century also were found in France (Bouchet and Paicheler, 1995).



FIGURE 1. *Schistosoma mansoni* egg with lateral spine (142 × 58 μm) in archaeological material (US 544) in Montbéliard site.

Schistosoma mansoni infection can only have originated from Africa because there are no intermediate hosts in Europe. A returning infected European voyager or a foreigner are 2 possibilities to be examined. Archaeological evidence showed that the house was occupied by a very rich family, which could have afforded a voyage to an epidemic region. However, there is no archaeological evidence that such a voyage was made. No souvenirs or exotic artifacts have been found. On the other hand, a slave could have introduced it. Since 1441, African slaves were taken to Portugal, and a few years after, they could be found in many European countries (Zurara, 1937). In France, there are records of African slaves by the middle of the 16th century (d'Avenel, 1910).

Together with the finding of *S. haematobium* (Bouchet and Paicheler, 1995) and *S. mansoni* eggs from the same archaeological material, evidence supports the presence of an infected human host, probably of African origin at this 15th–16th century European site. Nevertheless, wherever it came from and whichever host it came in, schistosomiasis was never established in Europe as an endemic disease.

The authors thank C. Goy (AFAN) for allowing us to study the archaeological material. Supported by CNRS, AFAN, CAPES/COFECUB, and CNPq/PRONEX.

LITERATURE CITED

- ARAÚJO, A. J. G., AND L. F. FERREIRA. 1997. Palaeoparasitology of schistosomiasis. *Memórias do Instituto Oswaldo Cruz* **92**: 717.
- D'AVENEL, V. 1910. *Découverte de l'histoire sociale*. Paris, France, 24 p.
- BOUCHET, E., AND J. C. PAICHELER. 1995. Palaeoparasitology: Presumption of a case of bilharzia on an archaeological site of the XVth century at Montbéliard (Doubs, France). *Comptes Rendus de l'Académie des Sciences de Paris (Ser III)* **318**: 811–814.
- BRUMPT, E. 1936. *Précis de Parasitologie*. Masson & Cie, Paris, France, 647 p.
- CANTRELLE, S., C. GOY, AND C. MUNIER. 2000. Histoire d'un quartier de Montbéliard (Doubs). D.A.F (Documents Archéologie française). *Maison des Sciences de L'Homme*, Paris, **83**, 140 p.
- CAPRON, L. 1993. 5000 ans de Bilharziose en Egypte. *Revue Pratique* **43**: 448.
- COCKBURN, A., R. A. BARRACO, T. A. REYMAN, AND W. H. PECK. 1975. Autopsy of an Egyptian mummy. *Science* **187**: 1155–1160.
- CONTIS, G., AND A. R. DAVID. 1996. The epidemiology of Bilharzia in ancient Egypt: 5000 years of schistosomiasis. *Parasitology Today* **12**: 253–255.
- DAVID, A. R. 1997. Disease in Egyptian mummies: The contribution of new technologies. *The Lancet* **349**: 1760–1763.
- DEELDER, A. M., R. L. MILLER, N. DE JONGE, AND F. W. KRIEGER. 1990. Detection of schistosome antigen in mummies. *The Lancet* **335**: 724–725.
- JONCKHEERE, F. 1942. *Autour de l'autopsie d'une momie*. Le Scribe royal Boute hamon. Fondation Egyptologique Reine Elisabeth, Bruxelles, Belgique, 137 p.
- LEWIN, P. K. 1977. Mummies that I have known. *American Journal of Diseases of Children* **131**: 349–350.
- LIANGBIAO, C., AND O. HUNG. 1981. Scanning electron microscopic view of parasitic worm ova. *Acta Academiologica Sinica* **3**: 1.
- MILLER, R. L., G. J. ARMELAGOS, S. IKRAM, N. DE JONGE, F. W. KRIEGER, AND A. M. DEELDER. 1992. Palaeoepidemiology of *Schistosoma* infection in mummies. *The British Medical Journal* **304**: 555–556.
- REINHARD, K., U. CONFALONIERI, B. HERRMANN, L. F. FERREIRA, AND A. ARAÚJO. 1988. Recovery of parasite remains from coprolites and latrines: Aspects of paleoparasitological technique. *Homo* **37**: 217–239.
- REYMAN, T. A. 1983. Les momies égyptiennes. *La Recherche* **14**: 792–799.
- ROLLISON, D., AND A. J. G. SIMPSON. 1987. The biology of schistosomes—From genes to latrines. Academic Press, London, U.K., 242 p.
- RUFFER, M. A. 1910. Note on the presence of *Bilharzia haematobium* in Egyptian mummies of the twentieth dynasty (1250–1000 BC). *The British Medical Journal* **1**: 16.
- WEI, O. 1973. Internal organs of a 2100-year-old female corpse. *The Lancet* **24**: 1198.
- ZURARA, G. E. 1937. *Crônica da Guiné*. Porto, Portugal, 108 p.